


FREDERICK COUNTY

RESULTS OF THE 1994-1997 MARYLAND BIOLOGICAL STREAM SURVEY: COUNTY ASSESSMENTS



**CHESAPEAKE BAY AND
WATERSHED PROGRAMS**
MONITORING AND
NON-TIDAL ASSESSMENT
CBWP-MANTA-EA-01-23





Parris N. Glendening
Governor

Kathleen Kennedy Townsend
Lt. Governor

A message to Maryland's citizens

The Maryland Department of Natural Resources (DNR) seeks to preserve, protect and enhance the living resources of the state. Working in partnership with the citizens of Maryland, this worthwhile goal will become a reality. This publication provides information that will increase your understanding of how DNR strives to reach that goal through its many diverse programs.

J. Charles Fox
Secretary

Karen M. White
Deputy Secretary



Maryland Department of Natural Resources
Tawes State Office Building
580 Taylor Avenue
Annapolis, Maryland 21401

Toll free in Maryland: 1-(877) 620 8DNR x8611
Out of state call: 410-260-8611
www.dnr.state.md.us

The facilities and services of the Maryland Department of Natural Resources are available to all without regard to race, color, religion, sex, sexual orientation, age, national origin, physical or mental disability.

This document is available in alternative format upon request from a qualified individual with a disability.

Publication date: December 2001
© MD DNR 2001



PRINTED ON RECYCLED PAPER

FREDERICK COUNTY

Results of the 1994-1997 Maryland Biological Stream Survey: County-Level Assessments

**Christopher J. Millard
Paul F. Kazyak
Anthony P. Prochaska**

December 2001

**Maryland Department of Natural Resources
Resource Assessment Service
Monitoring and Non-Tidal Assessment Division
580 Taylor Avenue
Annapolis, MD 21401**

**THIS PAGE INTENTIONALLY
LEFT BLANK**

FOREWORD

This report is based on results of the Maryland Biological Stream Survey (MBSS), a program funded primarily by the Power Plant Research Program and administered by the Maryland Department of Natural Resources (MDNR). Field data for the MBSS were collected by the Maryland Department of Natural Resources. Analyses of water chemistry samples were conducted by the University of Maryland's Appalachian Laboratory. Much of the initial data analysis was conducted by Versar, Inc. for MDNR's Power Plant Assessment Division.

This report helps fulfill two outcomes in MDNR's Strategic Plan: 1) A Vital and Life Sustaining Chesapeake Bay and Its Tributaries, and 2) Sustainable Populations of Living Resources and Healthy Ecosystems.

ACKNOWLEDGMENTS

The 1994-1997 Maryland Biological Stream Survey has been a cooperative effort among several agencies, consultants and academic institutions. We wish to thank Nancy Roth and Ginny Mercurio from Versar in helping to compile some of the data used in this report. Versar also designed the sampling program, obtained landowners' permissions, and helped manage the data. We are also grateful to the many individuals from Maryland Department of Natural Resources, the University of Maryland's Appalachian Laboratory (AL), and the University of Maryland's Wye Research and Education Center (WREC) who comprised the field crews and did a great job collecting the data. MDNR staff also digitized watersheds and calculated land use data, provided quality assurance, and conducted field crew training. Nancy Roth and her colleagues at Versar developed the fish Index of Biotic Integrity, and Dr. Sam Stribling and his staff at Tetra Tech, Inc. developed the benthic Index of Biotic Integrity. Dr. Ray Morgan of AL and Mr. Lenwood Hall of the WREC supervised additional field crews and developed the Physical Habitat Index, and Dr. Keith Eshleman of AL assisted with analyses of data on acidified streams. Drs. Wayne Starnes and Bob Reynolds of the Smithsonian Institution (reptiles and amphibians), Dr. Rich Raesly of Frostburg State University (fish), Rita Villella of the U.S. Geological Survey Leetown Science Center (mussels), and Michael Naylor of MDNR (aquatic vegetation) provided taxonomic verifications of voucher specimens. The success of the project resulted from the strong efforts of all these groups. Special thanks go to Ron Klauda for his editorial support and Brenda Morgan for her assistance in formatting, editing, and organizing the report.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Table of Contents

FOREWORD	i
ACKNOWLEDGEMENTS	i
INTRODUCTION	1
METHODS	3
MARYLAND BIOLOGICAL STREAM SURVEY DATA	9
COUNTY SUMMARY	10
LIST OF TABLES	
Table 1. Site information and land use data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	11
Table 2. Percent occurrence of fish species collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	15
Table 3. Tolerance Value, Functional Feeding Group, Habit, and Percent Occurrence of benthic macroinvertebrate taxa collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	17
Table 4. Percent occurrence of reptile and amphibian species collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	22
Table 5. Physical habitat data for Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	23
Table 6. Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	27
Table 7. Water chemistry data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	29
LIST OF FIGURES	
Figure 1. Land use in Frederick County.	13
Figure 2. Location of Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	14
Figure 3. Stream ecological conditions based on the Fish Index of Biotic Integrity (F-IBI) at Maryland Biological Stream Survey sites in Frederick County.	16
Figure 4. Stream ecological conditions based on the Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI) at Maryland Biological Stream Survey sites in Frederick County.	21
Figure 5. Stream ecological conditions based on the Physical Habitat Index (PHI) at Maryland Biological Stream Survey sites in Frederick County.	26
Figure 6. Nitrate-nitrogen concentrations (mg/L) at Maryland Biological Stream Survey sites in Frederick County.	31
LITERATURE CITED	33
APPENDIX A Summary of the types of data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.	A-1
APPENDIX B Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997.	B-1

**THIS PAGE INTENTIONALLY
LEFT BLANK**

INTRODUCTION

This report presents county-level data from the 1994–1997 Maryland Biological Stream Survey (MBSS or the Survey). Previous reports have documented interim results from the 1995 (Roth et al. 1997) and 1996 (Roth et al. 1998a) sample years. In addition, a comprehensive final report was produced to assess the “state of the streams” throughout the state (Roth et al. 1999). All previous MBSS reports have presented information by individual drainage basins. Because there is a recognized need for stream health information at the county level, a series of reports were prepared; this report is part of that series. This introductory section recounts the origin of the Survey and describes its components.

Origin of the MBSS

More than 10 years ago, the Maryland Department of Natural Resources (MDNR) recognized that atmospheric deposition was one of the most important environmental problems resulting from the generation of electric power. To determine the extent of acidification of Maryland streams resulting from acidic deposition, MDNR conducted the Maryland Synoptic Stream Chemistry Survey (MSSCS) in 1987. The MSSCS estimated the number and extent of streams at that time affected by or sensitive to acidification statewide and demonstrated the potential for adverse effects on biota from acidification. However, little direct information was available on the biological responses of Maryland streams to water chemistry conditions. Data that were available could not be used (because of methodological differences and spatial coverage limitations) to compare conditions across regions or watersheds (Tornatore et al. 1992). Neither was it possible to assess the interactions between acidic deposition and other anthropogenic and natural influences (CBRM 1989). For these reasons, in 1993, MDNR created the MBSS to provide comprehensive information on the status of biological resources in Maryland streams and how they are affected by acidic deposition and other cumulative effects of anthropogenic stresses.

Description of the MBSS

The MBSS is intended to help environmental decision-

makers protect and restore the natural resources of Maryland. The primary objectives of the MBSS are:

- to assess the current status of biological resources in Maryland’s non-tidal streams;
- to quantify the extent to which acidic deposition has affected or may be affecting biological resources in the state;
- to examine which other water chemistry, physical habitat, and land use factors are important in explaining the current status of biological resources in streams;
- to compile the first statewide inventory of stream biota;
- to establish a benchmark for long-term monitoring of trends in these biological resources; and
- to target future local-scale assessments and mitigation measures needed to restore degraded biological resources.

In creating the Survey, MDNR implemented a probability-based sampling design as a cost-effective way to characterize statewide stream resources. By randomly selecting sites, the Survey can make quantitative inferences about the characteristics of all 9,258 miles of first-to-third-order, non-tidal streams in Maryland (based on stream length on a 1:250,000-scale base map). MDNR recognized that the utility of these estimates depended on accurately measuring appropriate attributes of streams. The Survey focuses on biology for two reasons: (1) organisms themselves have direct societal value and (2) biological communities integrate stresses over time and are a valuable and cost-effective means of assessing ecological integrity (i.e., the capacity of a resource to sustain its inherent potential).

Fish are an important component of stream integrity and one that also contributes to substantial recreational values. For these reasons, fish communities are a primary focus of the Survey. The Survey collects quantitative data for the calculation of population estimates for individual fish species (both game and nongame). These data can also be used to evaluate

fish community composition, individual fish health, and the geographic distribution of commercially important, rare, or non-indigenous fish species. Benthic (bottom-dwelling) macroinvertebrates are another essential component of streams and they constitute the second principal focus of the Survey. The Survey uses rapid bioassessment procedures for collecting benthic macroinvertebrates; these semi-quantitative methods permit comparisons of relative abundance and community composition, and have proven to be an effective way of assessing biological integrity in streams (Hilsenhoff 1987, Lenat 1988, Plafkin et al. 1989, Kerans and Karr 1994, Resh 1995). The Survey also records the presence of reptiles and amphibians (herpetofauna), freshwater mussels, and aquatic plants (both submerged aquatic vegetation (SAV) and emergent macrophytes). The Survey has established rigorous protocols (Kazyak 1996) for each of these sampling components, as well as training and auditing procedures to assure that data quality objectives are met.

Although the MBSS sampling design and protocols provide exceptional information for characterizing the stream resources in Maryland, designation of degraded areas and identification of likely stresses requires additional activities. Assessing the condition of biological resources (whether they are degraded or not degraded) requires the development of ecological indicators that permit the comparison of sampled segment results to minimally impacted reference conditions (i.e., the biological community expected in watersheds with little or no human-induced impacts). The Survey has used its growing database of information collected with consistent methods and broad coverage across the state to develop and test indicators of individual biological components (Stribling et al. 1998, Roth et al. 1998b) and physical habitat quality (Hall et al. 1999). Each of these indicators consists of multiple metrics using the general approach developed for the Index of Biotic Integrity (IBI) (Karr et al. 1986, Karr 1991) and the Chesapeake Bay Benthic Restoration Goals (Ranasinghe et al. 1994). The fish and benthic macroinvertebrate IBIs (which combine attributes of both the number and the type of species found) are widely accepted indicators that have been adapted for use in a variety of geographic locations (Miller et al. 1988, Cairns and Pratt 1993, Simon 1999). The Survey is investigating the possibility

of developing additional indicators (e.g., amphibians in small streams with few or no fish) and combining components into a composite indicator of biological integrity.

In addition to developing reference-based indicators, the Survey is applying a variety of analytical methods to the question of which stressors are most closely associated with degraded streams. This involves correlational and multivariate analyses of water chemistry, physical habitat, land use, and biological information (e.g., presence of non-native species). The biological information also provides a valuable opportunity for documenting aquatic biodiversity across the state; the distribution and abundance of species previously designated as rare only by anecdotal evidence can be determined, and unique combinations of species at the ecosystem and landscape levels can be identified. Land use and other landscape-scale metrics will play an important role in identifying the relative contributions of different stressors to the cumulative impact on stream resources. Ultimately, the Survey seeks to provide an integrated assessment of the problems facing Maryland streams that will facilitate interdisciplinary solutions for their restoration. The survey also provides resource managers with the locations of relatively undisturbed streams and watersheds that deserve protection.

METHODS

This section presents the specific study design and procedures used to implement the Maryland Biological Stream Survey. The study area of concern and the sampling design developed to characterize it are presented, along with field and laboratory methods for each component: fish, benthic macroinvertebrates, reptiles and amphibians, physical habitat, and water chemistry. Methods for aquatic vegetation and mussel sampling are presented, but the resulting data are not included in this report. A full description of MBSS methods can be found in Kazyak (1996).

MBSS Study Design

The Survey study area comprises 17 distinct drainage basins across the state. Random sampling was used to allow the estimation of unbiased summary statistics (e.g., means, proportions, and their respective variances) for the entire state, a particular basin, and subpopulations of interest (e.g., streams with pH < 5).

Because it would have been cost prohibitive to visit a sufficient number of sites in all basins in a single year, lattice sampling was used to schedule sampling of all basins over a three-year period, 1995-1997. Lattice sampling, also known as multistratification, is a cost-effective means of allocating effort across time in a large geographic area (Heimbuch 1999, Jessen 1978, Cochran 1977). A table, or lattice, was formed by arranging 17 basins in 17 rows, and the years in 3 columns. Lattice sampling was the method used for selecting cells from this 17x3 table so that all basins would be sampled over a three-year period and all basins would have a non-zero probability of being sampled in a given year. The data presented in this report include those collected at random sampling sites within the 17 principal basins in Maryland, as well as sites from the 1994 demonstration project. Because no estimates were calculated for this report, these data were included to supplement the number of sites.

The sampling frame for the Survey was constructed by overlaying basin boundaries on a map of all blue-line stream reaches in the study area as digitized on a U.S. Geological Survey 1:250,000 scale topographic map. This sample frame was similar to that used by the earlier Maryland Synoptic Stream Chemistry Survey

(MSSCS) conducted in 1987 (Knapp and Saunders 1987, Knapp et al. 1988). The Strahler convention (Strahler 1957) was used for ranking stream reaches by order; first-order reaches, for example, are the most upstream reaches in the branching stream system. Sampling was restricted to non-tidal, third-order and smaller stream reaches, excluding impoundments that were non-wadable or that substantially altered the riverine nature of the reach (Kazyak 1994). Together, these first-through third-order streams comprise about 90% of all stream and river miles in Maryland. Stream reaches were further divided into non-overlapping, 75-meter segments; these segments were the elementary sampling units from which biological, water chemistry, and physical habitat data were collected.

The 1995-1997 MBSS study design was based on stratified random sampling of segments within each basin; each basin was stratified by stream order. Within a stream order, the number of segments sampled per basin is proportional to the number of stream miles in the basin. To achieve the target number of samples per stream order within each basin, a given number of segments were randomly selected from each basin and ranked in order of selection. In all basins, extra segments were selected as a contingency against loss of sampling sites from restricted access to selected streams or from streams that were dry, too deep, or otherwise unsampleable owing to field conditions. In some basins, where only a small number of sites would have been selected using this method, additional random sites were selected to increase sample size. These extra sites (selected at random using the method described above) were used to provide better basinwide estimates; they were not included in the estimates of statewide conditions.

Permissions were obtained to access privately owned land adjacent to or near each stream segment. The procedures for obtaining permissions are described in Chaillou (1995). Because landowner permissions were obtained in a synoptic fashion and some variation in these rates occurred, we obtained more permissions than were needed for the Survey. Only the highest ranking sites were sampled until the target goal for that basin was reached. For the three year study, the success rate for obtaining permission to access stream sampling segments was high. Eighty-eight percent of sites that were targeted for permission were sampled.

Reasons for permission denial varied and generally reflected the preferences of landowners regarding property access, rather than any specific types of land. In rare cases, permission denial may affect the interpretation of Survey estimates, but only where denials occur in streams with characteristics that differ from the general population of streams. In one example of potential bias, several sites with known coal mining activities in the North Branch Potomac basin denied permission to sample, likely under representing the proportion of acid mine drainage streams in the population.

Field and Laboratory Methods

Benthic macroinvertebrate and water quality sampling were conducted in spring, when the benthos are thought to be reliable indicators of environmental stress (Plafkin et al. 1989) and when acid deposition effects are often the most pronounced. Fish, reptiles and amphibians, aquatic vegetation, and mussel sampling, along with physical habitat evaluations, were conducted during the low-flow period in summer. Fish community composition tends to be stable during summer, and low flow is advantageous for electrofishing. Because low-flow conditions in summer may be a primary factor limiting the abundance and distribution of fish populations, habitat assessments were performed during the summer. The sample size in summer is lower than in spring because some streams were dry in summer or were, in rare cases, otherwise unsampleable.

To reduce temporal variability, sampling during spring and summer was conducted within specific, relatively narrow time intervals, referred to as index periods (Janicki et al. 1993). These index periods were defined by degree-day limits for specific parts of the state. This approach provided a synoptic assessment of the current status of stream biota, water quality, and physical habitat in the 17 basins sampled. The spring index period was the time period between approximately March 1 and May 1, with end of the index period determined by degree-day accumulation as specified in Hilsenhoff (1987). In reality, most spring samples (78%) were collected in March, well before degree-day accumulation limits were approached. The summer index period was between June 1 and September 30 (Kazyak 1994).

Data Collection and Measurement

Field sampling followed procedures specified in the MBSS sampling manual (e.g., Kazyak 1996). A summary of the variables measured and the field and laboratory methods used to conduct the sampling follows.

Fish

Fish were sampled during the summer index period using double-pass electrofishing within 75-meter stream segments. Block nets were placed at each end of the segment and direct current backpack electrofishing units were used to sample the entire segment. An attempt was made to thoroughly fish each segment, and consistent effort was applied over the two passes. This sampling approach allowed calculation of several metrics useful in calculating a biological index and produced unbiased estimates of fish species abundance.

In small streams, a single electrofishing unit was used. In larger streams, two to five units were employed to effectively sample the site. Captured fish were identified to species, counted, weighed, and released. Any individuals that could not be identified to species were retained for laboratory confirmation. For each pass, all individuals of each gamefish species (defined as trout, bass, walleye, pike, chain pickerel, and striped bass) were measured for total length and examined for visible external pathologies or anomalies. For nongame species, up to 100 fish of each species (from both passes) were examined for visible external pathologies or anomalies. For each pass, all non-game species were weighed together for an aggregate biomass measurement; gamefish were also weighed in aggregate to the nearest 10 g.

Electrofishing was also conducted at supplemental, non-randomly selected sites during the summer index period. The presence of each species of fish was recorded for these segments to provide additional qualitative information on statewide fish distributions. Sampling effort at most qualitative sites was based on doubling the elapsed time since the last species was recorded or a minimum of 600 seconds of electrofishing effort.

After processing the fish collected in the field, voucher

specimens were retained for each species not previously collected in the drainage basin. In addition, all individuals which could not be positively identified in the field were retained. The remaining fish were released. All voucher specimens and fish retained for positive identification in the laboratory were examined and verified by the MBSS Quality Assurance Officer or ichthyologists at Frostburg State University, Frostburg, Maryland or the Smithsonian Institution, Washington, DC.

Benthic Macroinvertebrates

Benthic macroinvertebrates were collected to provide a qualitative description of the community composition at each sampling site (Kazyak 1996). Sampling was conducted during the spring index period. Benthic community data were collected for the purpose of calculating biological metrics, such as those described in EPA's Rapid Bioassessment Protocols (Plafkin et al. 1989), and use as an indicator of biological integrity for Maryland streams.

At each segment, a 600 micron mesh "D" net was used to collect organisms from habitats likely to support the greatest taxonomic diversity. A riffle area was preferred, but other habitats were also sampled using a variety of techniques including kicking, jabbing, and gently rubbing hard surfaces by hand to dislodge organisms. If available, other habitat types were sampled, including rootwads, woody debris, leaf packs, macrophytes, and undercut banks. Each jab covered one square foot, and a total of approximately 2.0 m² (20 square feet) of combined substrates was sampled and preserved in 70% ethanol. In the laboratory, the preserved sample was transferred to a gridded pan and organisms were picked from randomly selected grid cells until the cell that contained the 100th individual (if possible) was completely picked. Some samples had fewer than 100 individuals. The benthic macroinvertebrates were identified to genus, or lowest practicable taxon, in the laboratory.

Index of Biotic Integrity

Sites were evaluated using both the fish (F-IBI) and benthic macroinvertebrate (B-IBI) IBIs developed for the MBSS (for detailed methods, see Roth et al. 1997 and Stribling et al. 1998). IBI scores for the MBSS are

determined by comparing the fish or benthic macroinvertebrate assemblages at each site to those found at minimally impacted reference sites. Three separate formulations were employed for the fish IBI, one for each of three distinct geographic areas: Coastal Plain, Eastern Piedmont, and Highland. The two formulations used for the benthic IBI cover the Coastal Plain and non-Coastal Plain regions. Individual metrics for the IBI are scored 1, 3, or 5, based on comparison with the distribution of metric values at reference sites. For either the individual metrics or total IBI, a score of 3 or greater is considered comparable to reference site conditions, while scores falling below this threshold differ significantly from the reference conditions. Scores for the MBSS IBIs are calculated as the mean of the individual metric scores and therefore range from 1 to 5. Some other programs have used a similar approach (e.g., Weisberg et al. 1997), while others have instead computed the IBI as the total of individual metric scores. For example, Karr et al. (1986) calculated IBI as the sum of 12 metric scores, with totals ranging from 12 to 60 points.

Reptiles and Amphibians

At each sample segment, reptiles and amphibians were identified and the presence of observed species was recorded during the summer index period. A search of the riparian area was conducted within 5 meters of the stream on both sides of the 75-meter segment. Any reptiles and amphibians collected during the electrofishing of the stream segment were also included in the species list. Individuals were identified to species when possible. Voucher specimens and individuals not positively identifiable in the field were retained for examination in the laboratory and confirmation by herpetologists at the Smithsonian Institution, Washington, DC, or Towson University, Towson, Maryland.

Physical Habitat

Habitat assessments were conducted at all stream segments as a means of assessing the importance of physical habitat to the biological integrity and fishability of freshwater streams in Maryland. Procedures for habitat assessments (Kazyak 1996) were derived from two currently used methodologies: EPA's Rapid

Bioassessment Protocols (RBPs) (Plafkin et al. 1989), as modified by Barbour and Stribling (1991), and the Ohio EPA's Qualitative Habitat Evaluation Index (QHEI) (Ohio EPA 1987, Rankin 1989). A number of characteristics (instream habitat, epifaunal substrate, velocity/depth diversity, pool/glide/eddy quality, riffle/run quality, channel alteration, bank stability, embeddedness, channel flow status, and shading) were assessed qualitatively, based on visual observations within each 75-meter sample segment. Riparian zone vegetation width was estimated to the nearest meter, up to 50 meters from the stream. Additional observations of the surrounding area were used to assign ratings for aesthetic value (based on visible signs of human refuse at a site) and remoteness (based on distance from the nearest road, accessibility, and evidence of human activity). Also recorded were the presence or absence of various stream features including substrate types, various morphological characteristics, beaver ponds, point sources, and stream channelization. Local land uses visible from the stream segment and riparian vegetation type were also noted. Several additional physical characteristics were measured quantitatively to further characterize the habitat for each segment (see Kazyak 1996 for details). Quantitative measurements of the segment included maximum depth, stream gradient, velocity, thalweg depth, number of functional rootwads, number of functional large woody debris, wetted width, sinuosity, and overbank flood height. A velocity/depth profile was measured or other data were collected to enable calculation of discharge.

Physical Habitat Index

The Physical Habitat Index (PHI) was developed using MBSS data from 1994 to 1997 (Hall et al. 1999). As was the case in development of the fish and benthic IBIs, the conceptual approach was based on evaluating the relative importance (discriminatory power) of individual metrics and combinations of metrics explaining natural differences in streams throughout Maryland. These metrics were derived from both quantitative and qualitative habitat data collected during the summer index period. Based on analyses conducted for both fish IBI (Roth et al. 1998) and benthic macroinvertebrate IBI (Stribling et al. 1998) development in Maryland, the State was divided into two regions: the Coastal Plain and non-Coastal Plain.

The resulting index was then adjusted to a centile scale that rated each sample segment as follows: Good - 72 to 100; Fair - 42 to 71.9; Poor - 12 to 41.9; and Very Poor - 0 to 11.9.

Water Chemistry

During the spring index period, water samples were collected at each site for analysis of pH, acid neutralizing capacity (ANC), conductivity, sulfate, nitrate-nitrogen, and dissolved organic carbon (DOC). These variables describe basic water quality conditions with an emphasis on factors related to acidic deposition.

Grab samples were collected in one-liter bottles for analysis of all analytes except pH. Water samples for pH were collected with 60 ml syringes, which allowed purging of air bubbles to minimize changes in carbon dioxide content (EPA 1987). Samples were stored on wet ice and shipped on wet ice to the analytical laboratory within 48 hours. Laboratory analyses were carried out by the University of Maryland's Appalachian Laboratory in Frostburg.

Chemical analysis of water samples followed standard methods described in EPA's Handbook of Methods for Acid Deposition Studies (EPA 1987). EPA protocols were followed, except that ANC sample volume was reduced to 40 ml to ease handling. Routine daily quality control (QC) checks included processing duplicate, blank, and calibration samples according to EPA guidelines for each analyte. Field duplicates were taken at 5% of all sites. Routine QC checks helped to identify and correct errors in sampling routines or instrumentation at the earliest possible stage.

During the summer index period, in situ measurements of dissolved oxygen (DO), pH, temperature, and conductivity were collected at each site to further characterize existing water quality conditions that might influence biological communities. Measurements were made at an undisturbed section of the segment, usually in the middle of the stream channel, using electrode probes. Instruments were calibrated daily and calibration logbooks were maintained to document instrument performance.

Recognizing that water temperature is an important factor affecting stream condition, but one that varies

daily and seasonally, temperature loggers were deployed at 220 sites in five basins during 1997. The basins sampled were: the Choptank, Susquehanna, Potomac Washington Metro, Patuxent, and Pocomoke. Onset Computer Corporation Optic Stowaway temperature loggers were anchored in each site during the summer index period. Water temperature was recorded every 15 minutes from June 15 until mid-September.

Mussels

During the summer index period, freshwater mussels were sampled qualitatively by examining each 75-meter stream segment for their presence. Mussels were identified to species, their presence recorded, and subsequently released. Species not positively identifiable in the field were retained for confirmation by U.S. Geological Survey (USGS) Biological Resources Division staff.

Aquatic Vegetation

Aquatic vegetation was sampled qualitatively by examining each 75-meter segment for the presence of aquatic plants. Plants were identified to species and their presence recorded for each site. While the primary objective was to document the presence of submerged aquatic vegetation (SAV), emergent and floating aquatic vegetation was also recorded when encountered. Species not positively identifiable in the field were retained for laboratory examination and confirmation by MDNR's staff expert on SAV. Due to the difficulty in long-term preservation, no permanent vouchers of aquatic vegetation were retained.

Data Management

All crews used standardized pre-printed data forms developed for the Survey to ensure that all data for each sampling segment were recorded and standard units of measure were used (Kazyak 1996). Using standard data forms facilitated data entry and minimized transcription error. The field crew leader and a second reviewer checked all data sheets for completeness and legibility before leaving each sampling location. Original data sheets were sent to the Data Management Officer for further review and data entry, while copies were retained by the field crews.

A custom database application, in which the input module was designed to match each of the field data sheets, was used for data entry. Data were independently entered into two databases and compared using a computer program as a quality-control procedure. Differences between the two databases were resolved from original data sheets or through discussions with field crew leaders.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Maryland Biological Stream Survey Data

COUNTY SUMMARY

A total of 74 quantitative sites were sampled in Frederick County by MBSS sampling crews during 1994-1997 (Table 1; Figure 2). Qualitative fish sampling was conducted at an additional ten sites to provide a more complete picture of fish species distributions. Appendix A provides a summary of the types of data available for each of the sites sampled.

Species Highlights

A total of 48 fish species were collected in the small to mid-sized streams that were sampled (Table 2); this number ties the county for a ranking of fourth in the state for species richness. Blacknose dace and fantail darter, two pollution-tolerant species, were the most commonly found fish species. In contrast, pollution-sensitive brook trout, once found in streams throughout Frederick County, were only collected at 4% of the streams sampled by the MBSS. No state or federally listed fish species were collected. However, pearl dace and checkered sculpin, two fish species rare in the state of Maryland, were each collected at a single site in Frederick County.

The 172 genera of benthic macroinvertebrates found ties this county for a ranking of ninth in the state, but nearly one-third of these taxa were found at a single site and many appear to be rare on a statewide basis (Table 3). In addition, only three of the genera collected could be found at more than half of the sites sampled.

Nineteen species of reptiles and amphibians were found in or near Frederick County streams (Table 4), ranking it eighth in the state. No state or federally listed reptiles or amphibians were collected during the sampling.

Ecological Health

The overall ecological health of county streams can best be described as Fair to Poor. The average F-IBI score among sites was 3.25 (Fair range, but low in the category), and the average B-IBI score was 2.30 (Poor range). Based on F-IBI and B-IBI scores, the highest rated small to mid-sized streams are South Fork Linganore Creek, Ben's Branch, Bush Creek, Ballenger Creek, Little Hunting Creek, and Fishing Creek (Table

6). Some of the lowest rated streams include: Addison Run, Clemson Branch, Sam's Creek, Motter's Run, Tuscarora Creek, Beaver Dam Branch, and an unnamed tributary to Carroll Creek.

Physical Habitat

Physical habitat in Frederick County was rated as Fair by the Physical Habitat Index. Values ranged from 3.55 to 99.93, with an average score of 53.22 (low end of the Fair range, ranking twelfth among counties in the state) (Table 6; Figure 5). Other noteworthy points include rankigns of seventh and sixth for epifaunal substrate and instream habitat, respectively. In addition, the county ranked twentieth for large woody debris (instream logs) abundance and fourteenth for instream rootwads (trees whose roots protect banks from erosion and provide habitat for aquatic life).

Nitrate-Nitrogen

Nitrate-nitrogen values at sites sampled were generally elevated and averaged 3.0 mg/L, making Frederick the ninth worst county in the state. In no stream was the EPA limit for drinking water (10 mg/L) exceeded (Table 7).

Table 1. Site information and land use data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997. Basin abbreviations are as follows: MP - Middle Potomac River.

Site	Latitude	Longitude	Stream Name	Basin	Order	Catchment Acres	% Urban	% Agric.	% Forest
CR-P-021-329-96	39.5480	77.1699	Sam's Cr	MP	3	10745.17	0.11	77.99	21.06
FR-B-032-206-96	39.7007	77.4578	Friends Cr	MP	2	1295.98	0.55	29.49	69.83
FR-B-046-127-96	39.5066	77.5124	Un Trib To Little Catoctin Cr	MP	1	300.43	0.11	0.34	99.55
FR-B-065-111-96	39.4494	77.5481	Un Trib To Catoctin Cr	MP	1	279.68	8.28	87.21	4.51
FR-B-076-118-96	39.3475	77.6182	Un Trib To Little Catoctin Cr	MP	1	209.25	4.91	80.03	12.93
FR-B-081-229-96	39.6295	77.4619	Hunting Cr	MP	2	2425.24	0.00	19.85	78.37
FR-B-085-212-96	39.4893	77.5711	Catoctin Cr	MP	2	22939.43	0.45	39.65	59.38
FR-B-133-222-96	39.6585	77.4842	Owens Cr	MP	2	1230.31	0.03	16.87	82.55
FR-B-164-137-96	39.6207	77.5280	Middle Cr	MP	1	760.78	0.04	8.42	91.18
FR-P-005-141-96	39.4398	77.4919	Un Trib To Carroll Cr	MP	1	553.57	3.94	26.32	69.68
FR-P-009-341-96	39.7076	77.3206	Flat Run	MP	3	6701.52	1.51	63.92	33.21
FR-P-009-347-96	39.7101	77.3199	Flat Run	MP	3	6070.93	0.56	63.53	34.44
FR-P-015-304-96	39.3183	77.3334	Bennet Cr	MP	3	16582.01	1.71	70.89	25.20
FR-P-034-228-96	39.5788	77.4310	Little Hunting Cr	MP	2	4874.64	0.13	1.30	98.26
FR-P-038-139-96	39.7011	77.3649	Turkey Cr	MP	1	1443.65	0.02	0.62	99.29
FR-P-046-227-96	39.5566	77.1816	Haines Br	MP	2	1967.28	0.14	87.93	10.19
FR-P-050-354-96	39.4915	77.3191	Cabbage Run	MP	3	4449.00	0.40	58.94	38.25
FR-P-093-237-96	39.4940	77.4355	Tuscarora Cr	MP	2	7690.82	0.06	9.65	90.17
FR-P-093-238-96	39.4936	77.4348	Tuscarora Cr	MP	2	7710.55	0.06	9.83	89.99
FR-P-100-117-96	39.3943	77.4815	Un Trib To Ballenger Cr	MP	1	434.49	0.47	39.13	59.37
FR-P-101-233-96	39.3267	77.3150	Fahrney Br	MP	2	5018.22	2.02	80.21	14.79
FR-P-103-230-96	39.3791	77.4604	Ballenger Cr	MP	2	2598.42	4.19	63.92	30.74
FR-P-111-134-96	39.6603	77.2737	Un Trib To Piney Cr	MP	1	315.40	0.00	94.70	5.30
FR-P-116-221-96	39.4984	77.2922	Un Trib To Israel Cr	MP	2	2137.74	0.21	55.64	43.68
FR-P-132-320-96	39.5661	77.3986	Little Hunting Cr	MP	3	7120.92	0.24	15.17	83.93
FR-P-156-217-96	39.4264	77.1753	South Fork Linganore Cr	MP	2	4372.10	0.24	65.30	32.63
FR-P-156-231-96	39.4277	77.1772	South Fork Linganore Cr	MP	2	4570.12	0.22	64.18	33.72
FR-P-156-234-96	39.4285	77.1903	South Fork Linganore Cr	MP	2	4864.43	0.20	63.25	34.60
FR-P-156-252-96	39.4238	77.1727	South Fork Linganore	MP	2	4104.67	0.25	65.88	32.08
FR-P-168-218-96	39.3264	77.4707	Tuscarora Cr	MP	2	2685.53	7.98	73.51	17.15
FR-P-214-303-96	39.5911	77.3371	Owens Cr	MP	3	25139.77	0.40	44.73	53.40
FR-P-214-342-96	39.5887	77.3392	Owens Cr	MP	3	25476.95	0.39	45.18	52.89
FR-P-223-225-96	39.5639	77.2111	Beaver Dam Cr	MP	2	3929.79	0.16	82.61	16.75
FR-P-223-240-96	39.5593	77.2097	Beaver Dam Br	MP	2	3568.79	0.18	83.85	15.46
FR-P-258-202-96	39.5070	77.3839	Fishing Cr	MP	2	5107.03	0.53	74.87	22.23
FR-P-258-215-96	39.5192	77.3875	Fishing Cr	MP	2	4628.99	0.59	74.44	22.51
FR-P-258-243-96	39.5110	77.3823	Fishing Cr	MP	2	5047.48	0.53	74.80	22.32
FR-P-261-122-96	39.5961	77.3910	Un Trib To Hunting Cr	MP	1	716.55	2.95	81.99	15.07
FR-P-263-311-96	39.2800	77.4873	Tuscarora Cr	MP	3	10275.64	2.95	70.31	25.54
FR-P-265-335-96	39.7156	77.2925	Middle Cr	MP	3	16059.29	0.93	40.80	56.27
FR-P-265-351-96	39.7176	77.2926	Middle Cr	MP	3	16000.20	0.94	40.81	56.26
FR-P-275-239-96	39.3565	77.2618	Church Br Of Bush Cr	MP	2	3109.13	4.58	64.48	29.64
FR-P-277-115-96	39.3866	77.4773	Un Trib To Ballenger Cr	MP	1	662.45	1.75	57.12	40.98
FR-P-288-133-96	39.6097	77.4156	Trib To Hunting Cr	MP	1	1036.38	4.34	6.90	88.76
FR-P-290-121-96	39.2975	77.5033	Un Trib To Tuscarora Cr	MP	1	439.93	2.86	53.67	41.61
FR-P-294-313-96	39.5707	77.3914	Hunting Cr	MP	3	14753.21	4.59	29.45	64.30
FR-P-294-357-96	39.5745	77.3932	Hunting Cr	MP	3	14497.01	4.65	28.40	65.25
FR-P-298-308-96	39.6084	77.3358	Owens Cr	MP	3	19411.55	0.24	39.06	59.25
FR-P-300-130-96	39.5830	77.3564	Un Trib To Monocacy R	MP	1	831.55	0.12	95.32	4.31

Table 1 (cont.). Site information and land use data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997. Basin abbreviations are as follows: MP - Middle Potomac River.

Site	Latitude	Longitude	Stream Name	Basin	Order	Catchment Acres	% Urban	% Agric.	% Forest
FR-P-302-334-96	39.6379	77.3848	Owens Cr	MP	3	9961.25	0.23	17.95	80.93
FR-P-319-352-96	39.6133	77.3591	Owens Cr	MP	3	16947.09	0.23	32.00	66.26
FR-P-321-214-96	39.4279	77.2012	Woodville Br	MP	2	4425.86	2.43	70.38	25.26
FR-P-335-110-96	39.4421	77.3971	Un Trib To Monocacy R	MP	1	501.52	21.76	73.12	5.12
FR-P-349-204-96	39.3847	77.4705	Ballenger Cr	MP	2	3531.03	3.52	53.54	41.77
FR-P-351-112-96	39.3419	77.2789	Un Trib To Fahrney Br	MP	1	173.18	0.00	95.27	2.17
FR-P-354-321-96	39.2537	77.4830	Tuscarora Cr	MP	3	12773.35	2.40	73.77	22.39
FR-P-360-220-96	39.3511	77.3654	Un Trib To Bush Cr	MP	2	844.48	2.51	83.98	12.34
FR-P-371-132-96	39.6535	77.3161	Motter's Run	MP	1	758.74	0.00	93.90	6.06
FR-P-377-242-96	39.3029	77.2796	Bennett Cr	MP	2	6294.81	1.50	62.92	33.43
FR-P-388-208-96	39.4542	77.1547	Talbot Br	MP	2	3038.36	1.12	57.80	39.11
FR-P-388-246-96	39.4511	77.1370	Talbot Br	MP	2	2740.65	1.23	57.09	39.96
FR-P-394-317-96	39.4933	77.3112	Cabbage Run	MP	3	4091.40	0.44	57.68	39.41
FR-P-399-126-96	39.4829	77.1505	Un Trib To Weldon Cr	MP	1	227.96	0.00	73.43	26.57
FR-P-409-210-96	39.4409	77.3504	Addison Run	MP	2	1584.51	0.17	63.58	35.90
FR-P-411-305-96	39.4198	77.2539	Bens Br	MP	3	6392.80	0.81	67.85	29.89
FR-P-421-306-96	39.3588	77.3667	Bush Cr	MP	3	19046.04	3.57	63.25	31.14
FR-P-429-307-96	39.3724	77.4472	Ballenger Cr	MP	3	7606.78	4.81	61.58	32.27
FR-P-461-251-96	39.4134	77.2371	Ben's Br	MP	2	4226.48	1.03	61.15	36.15
FR-P-462-346-96	39.5197	77.1710	Clemson Br	MP	3	2130.59	0.14	85.31	14.33
FR-P-474-302-96	39.5651	77.1860	Sam's Cr	MP	3	14058.44	0.43	79.08	19.03
FR-P-479-348-96	39.6445	77.4147	Owens Cr	MP	3	8418.60	0.16	12.80	86.45
FR-P-516-235-96	39.5570	77.3945	Sandy Run	MP	2	2208.85	1.22	63.93	34.41
FR-P-545-325-96	39.3610	77.3083	Bush Cr	MP	3	11915.60	5.36	59.20	32.86
FR-P-545-345-96	39.3602	77.3096	Bush Cr	MP	3	11939.23	5.37	59.16	32.87




Figure 1. Land use in Frederick County (MOP 1994).




Figure 2. Location of Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Table 2. Percent occurrence of fish species collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Family	Common Name	Scientific Name	Number of Occurrences	Percent Occurrence
Anguillidae	American eel	<i>Anguilla rostrata</i>	3	4.11
Cyprinidae	central stoneroller	<i>Camptostoma anomalum</i>	36	49.32
	rosyside dace	<i>Clinostomus funduloides</i>	19	26.03
	satinfin shiner	<i>Cyprinella analostana</i>	3	4.11
	spotfin shiner	<i>Cyprinella spiloptera</i>	19	26.03
	common carp	<i>Cyprinus carpio</i>	1	1.37
	cutlips minnow	<i>Exoglossum maxillingua</i>	14	19.18
	common shiner	<i>Luxilus cornutus</i>	25	34.25
	pearl dace	<i>Margariscus margarita</i>	1	1.37
	river chub	<i>Nocomis micropogon</i>	7	9.59
	golden shiner	<i>Notemigonus crysoleucas</i>	2	2.74
	silverjaw minnow	<i>Notropis buccatus</i>	14	19.18
	spottail shiner	<i>Notropis hudsonius</i>	11	15.07
	swallowtail shiner	<i>Notropis procne</i>	1	1.37
	rosyface shiner	<i>Notropis rubellus</i>	1	1.37
	bluntnose minnow	<i>Pimephales notatus</i>	44	60.27
	fathead minnow	<i>Pimephales promelas</i>	1	1.37
	blacknose dace	<i>Rhinichthys atratulus</i>	69	94.52
	longnose dace	<i>Rhinichthys cataractae</i>	53	72.60
	creek chub	<i>Semotilus atromaculatus</i>	54	73.97
	fallfish	<i>Semotilus corporalis</i>	25	34.25
Catostomidae	white sucker	<i>Catostomus commersoni</i>	50	68.49
	creek chubsucker	<i>Erimyzon oblongus</i>	3	4.11
	northern hogsucker	<i>Hypentelium nigricans</i>	19	26.03
	golden redhorse ¹	<i>Moxostoma erythrurum</i>		
Ictaluridae	yellow bullhead	<i>Ameiurus natalis</i>	25	34.25
	brown bullhead	<i>Ameiurus nebulosus</i>	1	1.37
	channel catfish	<i>Ictalurus punctatus</i>	1	1.37
	marginated madtom	<i>Noturus insignis</i>	6	8.22
Salmonidae	rainbow trout	<i>Oncorhynchus mykiss</i>	1	1.37
	brown trout	<i>Salmo trutta</i>	5	6.85
	brook trout	<i>Salvelinus fontinalis</i>	3	4.11
Cyprinodontidae	banded killifish	<i>Fundulus diaphanus</i>	11	15.07
Cottidae	mottled sculpin	<i>Cottus bairdi</i>	32	43.84
	checkered sculpin	<i>Cottus sp.</i>	1	1.37
	Potomac sculpin	<i>Cottus girardi</i>	44	60.27
Percichthyidae	white perch ¹	<i>Morone americana</i>		
Centrarchidae	rock bass	<i>Ambloplites rupestris</i>	23	31.51
	redbreast sunfish	<i>Lepomis auritus</i>	34	46.58
	green sunfish	<i>Lepomis cyanellus</i>	29	39.73
	pumpkinseed	<i>Lepomis gibbosus</i>	13	17.81
	bluegill	<i>Lepomis macrochirus</i>	43	58.90
	longear sunfish	<i>Lepomis megalotis</i>	2	2.74
	smallmouth bass	<i>Micropterus dolomieu</i>	13	17.81
	largemouth bass	<i>Micropterus salmoides</i>	25	34.25
Percidae	greenside darter	<i>Etheostoma blennioides</i>	28	38.36
	fantail darter	<i>Etheostoma flabellare</i>	58	79.45
	tessellated darter	<i>Etheostoma olmstedii</i>	8	10.96
None			1	1.37

¹ Qualitative Sites




Figure 3. Stream ecological conditions based on the Fish Index of Biotic Integrity (F-IBI) at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Table 3. Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - clinger, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Turbellaria	Tricladida	Planariidae	<i>Cura</i> Sp.			sp	2.70
			<i>Dugesia</i> Sp.	7	Predator	sp	2.70
Oligochaeta	Lumbriculida	Lumbriculidae		10	Collector	bu	5.41
Oligochaeta	Tubificida	Enchytraeidae		10	Collector	bu	4.05
		Naididae		10	Collector	bu	5.41
		Tubificidae		10	Collector	cn	12.16
			<i>Limnodrilus</i> Sp.	10	Collector	cn	6.76
Hirudinea				8	Predator	sp	2.70
Gastropoda	Basommatophora	Ancylidae	<i>Fissia</i> Sp.	7	Scraper	cb	1.35
		Lymnaeidae	<i>Pseudosuccinea</i> Sp.	6	Collector	cb	2.70
		Physidae	<i>Physella</i> Sp.	8	Scraper	cb	6.76
		Planorbidae	<i>Helisoma</i> Sp.	6	Scraper	cb	1.35
Pelecypoda	Veneroida	Corbiculidae	<i>Corbicula</i> Sp.	6	Filterer	bu	1.35
		Sphaeriidae			Filterer	bu	1.35
			<i>Pisidium</i> Sp.	8	Filterer	bu	6.76
			<i>Sphaerium</i> Sp.	8	Filterer	bu	6.76
Malacostraca	Amphipoda	Crangonyctidae	<i>Crangonyx</i> Sp.	4	Collector	sp	8.11
		Gammaridae	<i>Gammarus</i> Sp.	6	Shredder	sp	5.41
			<i>Stygonectes</i> Sp.	6	Shredder	sp	2.70
			<i>Hyalella</i> Sp.	6	Shredder	sp	2.70
Malacostraca	Decapoda	Cambaridae		6	Shredder	sp	2.70
			<i>Cambarus</i> Sp.	6	Collector	sp	1.35
Malacostraca	Isopoda	Asellidae	<i>Caecidotea</i> Sp.	8	Collector	sp	4.05
			<i>Lirceus</i> Sp.	8	Collector	sp	2.70
Insecta	Ephemeroptera	Ameletidae	<i>Ameletus</i> Sp.	0	Collector	sw, cb	8.11
		Baetidae			Collector	sw, cn	10.81
			<i>Acerpenna</i> Sp.	4	Collector	sw, cn	6.76
			<i>Baetis</i> Sp.	6	Collector	sw, cb, cn	6.76
			<i>Centroptilum</i> Sp.	2	Collector	sw, cn	1.35
			<i>Dipheter</i> Sp.		Collector	sw, cn	1.35
		Baetiscidae	<i>Baetisca</i> Sp.	4	Collector	sp	1.35
		Caenidae	<i>Caenis</i> Sp.	7	Collector	sp	2.70
		Ephemerellidae				cn, sp, sw	4.05
			<i>Ephemerella</i> Sp.	2	Collector	cn, sw	33.78
			<i>Eurylophella</i> Sp.	4	Scraper	cn, sp	18.92
			<i>Satella</i> Sp.	2	Collector	cn	1.35
			<i>Ephemera</i> Sp.	3	Collector	bu	5.41
					Scraper	cn	2.70
			<i>Cinygmula</i> Sp.		Scraper	cn	1.35
			<i>Epeorus</i> Sp.	0	Scraper	cn	8.11
			<i>Leucrocota</i> Sp.	1	Scraper	cn	2.70
			<i>Stenacron</i> Sp.	4	Collector	cn	4.05
			<i>Stenonema</i> Sp.	4	Scraper	cn	39.19
		Isonychiidae	<i>Isonychia</i> Sp.	2	Filterer	sw, cn	21.62
		Leptophlebiidae			Collector	sw, cn	5.41
			<i>Leptophlebia</i> Sp.	4	Collector	sw, cn, sp	4.05
			<i>Paraleptophlebia</i> Sp.	2	Collector	sw, cn, sp	18.92

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - climber, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Insecta	Odonata	Potamanthidae	<i>Anthopotamus</i> Sp.	4			1.35
		Aeshnidae	<i>Boyeria</i> Sp.	2	Predator	cb, sp	1.35
		Calopterygidae	<i>Calopteryx</i> Sp.	6	Predator	cb	1.35
		Coenagrionidae	<i>Argia</i> Sp.	8	Predator	cn, cb, sp	2.70
		Corduliidae	<i>Macromia</i> Sp.	3	Predator	sp	1.35
		Gomphidae			Predator	bu	1.35
Insecta	Plecoptera		<i>Lanthus</i> Sp.	6	Predator	bu	1.35
			<i>Stylogomphus</i> Sp.		Predator	bu	6.76
		Capniidae			Shredder	sp, cn	6.76
			<i>Allocapnia</i> Sp.	3	Shredder	cn	21.62
			<i>Paracaenia</i> Sp.	1	Shredder	-	27.03
		Chloroperlidae	<i>Sweltsa</i> Sp.		Predator	cn	5.41
		Leuctridae	<i>Leuctra</i> Sp.	0	Shredder	cn	4.05
		Nemouridae			Shredder	sp, cn	2.70
			<i>Amphinemura</i> Sp.	3	Shredder	sp, cn	22.97
			<i>Ostrocerca</i> Sp.		Shredder	sp, cn	1.35
			<i>Prostoia</i> Sp.		Shredder	sp, cn	25.68
		Peltoperlidae	<i>Tallaperla</i> Sp.		Shredder	cn, sp	1.35
		Perlidae	<i>Acronemura</i> Sp.	0	Predator	cn	10.81
			<i>Paragnetina</i> Sp.	1	Predator	cn	1.35
		Perlodidae			Predator	cn	4.05
			<i>Clasperla</i> Sp.	1	Predator	cn	2.70
			<i>Isoperla</i> Sp.	2	Predator	cn, sp	1.35
		Pteronarcyidae	<i>Pteronarcys</i> Sp.	2	Shredder	cn, sp	2.70
		Taeniopterygidae		2	Shredder		2.70
Insecta	Hemiptera		<i>Oemopteryx</i> Sp.		Shredder	sp, cn	10.81
			<i>Strophopteryx</i> Sp.		Shredder	sp, cn	20.27
Insecta	Megalopectera	Corixidae			Predator	sw	2.70
Insecta		Corydalidae	<i>Corydalus</i> Sp.	5	Predator	cn, cb	2.70
			<i>Nigronia</i> Sp.	0	Predator	cn, cb	9.46
		Sialidae	<i>Sialis</i> Sp.	4	Predator	bu, cb, cn	5.41
Insecta	Trichoptera	Hydropsychidae	<i>Cheumatopsyche</i> Sp.	5	Filterer	cn	48.65
			<i>Diplectrona</i> Sp.	2	Filterer	cn	18.92
			<i>Hydropsyche</i> Sp.	6	Filterer	cn	31.08
		Hydroptilidae	<i>Hydroptila</i> Sp.	6	Scraper	cn	1.35
		Lepidostomatidae	<i>Lepidostoma</i> Sp.	3	Shredder	cb, sp, cn	1.35
		Leptoceridae	<i>Mystacides</i> Sp.	4	Collector	sp, cb	1.35
		Limnephilidae			Shredder	cb, sp, cn	2.70
			<i>Pycnopsyche</i> Sp.	4	Shredder	sp, cb, cn	1.35
		Philopotamidae	<i>Chimarra</i> Sp.	4	Filterer	cn	22.97
			<i>Dolophilodes</i> Sp.	0	Filterer	cn	1.35
		Polycentropodidae	<i>Polycentropus</i> Sp.	5	Filterer	cn	1.35
		Rhyacophilidae	<i>Rhyacophila</i> Sp.	1	Predator	cn	14.86
		Uenoidae	<i>Neophylax</i> Sp.	3	Scraper	cn	24.32
Insecta	Coleoptera	Dytiscidae	<i>Agabus</i> Sp.	5	Predator	sw, dv	1.35
			<i>Hydroporus</i> Sp.	5	Predator	sw, cb	1.35
			<i>Ancyronyx</i> Sp.	2	Scraper	cn, sp	2.70

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - clinger, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
Insecta	Diptera	Halipilidae	<i>Dubiraphia</i> Sp.	6	Scraper	cn, cb	12.16
			<i>Macronychus</i> Sp.	4	Scraper	cn	6.76
			<i>Optioservus</i> Sp.	4	Scraper	cn	47.30
			<i>Oulimnius</i> Sp.	2	Scraper	cn	14.86
			<i>Promoresia</i> Sp.	2	Scraper	cn	1.35
			<i>Stenelmis</i> Sp.	6	Scraper	cn	35.14
			<i>Pelodytes</i> Sp.	5	Shredder	cb, cn	1.35
		Hydrophilidae	<i>Hydrobius</i> Sp.	5	Collector	cb, cn, sp	1.35
		Psephenidae	<i>Ectopria</i> Sp.	5	Scraper	cn	2.70
		Psephenidae	<i>Psephenus</i> Sp.	4	Scraper	cn	8.11
			<i>Anchytarsus</i> Sp.	4	Shredder	cn	1.35
		Ceratopogonidae			Predator	sp, bu	1.35
			<i>Bezzia</i> Sp.	6	Predator	bu	4.05
		Chironomidae	<i>Ceratopogon</i> Sp.	6	Predator	sp, bu	1.35
			<i>Helius</i> Sp.	4	Predator	sp, bu	1.35
		Chironomidae					2.70
			<i>Brillia</i> Sp.	5	Shredder	bu, sp	6.76
			<i>Chaetocladius</i> Sp.	6	Collector	sp	1.35
			<i>Chironomus</i> Sp.	10	Collector	bu	9.46
			<i>Conchapelopia</i> Sp.	6	Predator	sp	27.03
			<i>Corynoneura</i> Sp.	7	Collector	sp	10.81
			<i>Cricotopus</i> Sp.	7	Shredder	cn, bu	4.05
			<i>Cricotopus</i> /				
			<i>Orthocladius</i> Sp.		Shredder		51.35
			<i>Cryptochironomus</i> Sp.	8	Predator	sp, bu	2.70
			<i>Diamesa</i> Sp.	5	Collector	sp	12.16
			<i>Dicortendipes</i> Sp.	10	Collector	bu	4.05
			<i>Diplocladius</i> Sp.	7	Collector	sp	1.35
			<i>Endochironomus</i> Sp.	10	Shredder	cn	1.35
			<i>Eukiefferiella</i> Sp.	8	Collector	sp	36.49
			<i>Heleniella</i> Sp.		Predator	sp	1.35
			<i>Heterotrissocladius</i> Sp.		Collector	sp, bu	6.76
			<i>Hydrobaenus</i> Sp.	8	Scraper	sp	8.11
			<i>Micropsectra</i> Sp.	7	Collector	cb, sp	9.46
			<i>Microtendipes</i> Sp.	6	Filterer	cn	18.92
			<i>Nanocladius</i> Sp.	3	Collector	sp	1.35
			<i>Orthoclaudiinae A</i> Sp.		Collector		12.16
			<i>Orthocladius</i> Sp.	6	Collector	sp, bu	25.68
			<i>Parachaetocladius</i> Sp.	2	Collector	sp	1.35
			<i>Parakiefferiella</i> Sp.	4	Collector	sp	2.70
			<i>Paramerina</i> Sp.	4	Predator	sp	4.05
			<i>Parametriocnemus</i> Sp.	5	Collector	sp	55.41
			<i>Paratanytarsus</i> Sp.	6	Collector	sp	2.70
			<i>Phaenopsectra</i> Sp.	7	Collector	cn	2.70
			<i>Polypedilum</i> Sp.	6	Shredder	cb, cn	10.81
			<i>Potthastia</i> Sp.	2	Collector	sp	1.35
			<i>Procladius</i> Sp.	9	Predator	sp	1.35

Table 3 (cont.). Tolerance Value (TV)¹, Functional Feeding Group (FFG), Habit, and Percent Occurrence of benthic macroinvertebrate taxa² collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997. Abbreviations of habits are as follows: bu - burrower, cn - climber, cb - climber, sp - sprawler, dv - diver, and sk - skater.

Class	Order	Family	Genus	TV	FFG	Habit	Percent Occurrence
			<i>Rheocricotopus</i> Sp.	6	Collector	sp	2.70
			<i>Rheotanytarsus</i> Sp.	6	Filterer	cn	10.81
			<i>Stempellinella</i> Sp.	4	Collector	cb, sp, cn	2.70
			<i>Stictochironomus</i> Sp.	9	Collector	bu	8.11
			<i>Sympotthastia</i> Sp.	2	Collector	sp	33.78
			<i>Tanytarsus</i> Sp.	6	Filterer	cb, cn	10.81
			<i>Thienemanniella</i> Sp.	6	Collector	sp	22.97
			<i>Thienemannimyia</i> Sp.		Predator	sp	17.57
			<i>Trissopelopia</i> Sp.		Predator	sp	1.35
			<i>Tvetenia</i> Sp.	5	Collector	sp	1.35
			DIAMESINAE		Collector		1.35
			ORTHOCLADIINAE		Collector		1.35
			TANYPODINAE		Predator		1.35
			TANYTARSINI		Collector		1.35
			<i>Zavrelimyia</i> Sp.	8	Predator	sp	12.16
		Dixidae	<i>Dixa</i> Sp.	4	Predator	sw, cb	1.35
		Empididae	<i>Chelifera</i> Sp.		Predator	sp, bu	1.35
			<i>Clinocera</i> Sp.		Predator	cn	13.51
			<i>Hemerodromia</i> Sp.	6	Predator	sp, bu	18.92
		Simuliidae		7	Filterer	cn	1.35
			<i>Prosimulium</i> Sp.	7	Filterer	cn	77.03
			<i>Simulium</i> Sp.	7	Filterer	cn	14.86
			<i>Stegopterna</i> Sp.	7	Filterer	cn	31.08
		Tabanidae	<i>Chrysops</i> Sp.	7	Predator	sp, bu	5.41
		Tipulidae			Predator	bu, sp	2.70
			<i>Antocha</i> Sp.	5	Collector	cn	10.81
			<i>Dicranota</i> Sp.	4	Predator	sp, bu	8.11
			<i>Hexatoma</i> Sp.	4	Predator	bu, sp	12.16
			<i>Limonia</i> Sp.	6	Shredder	bu, sp	2.70
			<i>Ormosia</i> Sp.		Collector	bu	4.05
			<i>Pseudolimnophila</i> Sp.	2	Predator	bu	10.81
			<i>Tipula</i> Sp.	4	Shredder	bu	24.32

¹ Tolerance values are on a 0 (extremely sensitive) to 10 (tolerant) scale.

² Taxa not identified to genus are presented in capital letters. Subfamily - Tanypodinae, Orthocladiinae, Diamesinae; Tribe - Tanytarsini.




Figure 4. Stream ecological conditions based on the Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI) at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Table 4. Percent occurrence of reptile and amphibian species collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Family	Common Name	Scientific Name	Number of Occurrences	Percent Occurrence
Plethodontidae	longtail salamander	<i>Eurycea l. longicauda</i>	3	4.11
	northern dusky salamander	<i>Desmognathus f. fuscus</i>	3	4.11
	northern two-lined salamander	<i>Eurycea bislineata</i>	28	38.36
	northern slimy salamander	<i>Plethodon glutinosus</i>	1	1.37
	northern spring salamander	<i>Gyrinophilus p. porphyriticus</i>	1	1.37
	red salamander	<i>Pseudotriton ruber</i>	7	9.59
	redback salamander	<i>Plethodon cinereus</i>	7	9.59
Bufonidae	American toad	<i>Bufo americanus</i>	11	15.07
	Fowler's toad	<i>Bufo woodhousii fowleri</i>	1	1.37
Hylidae	northern spring peeper	<i>Pseudacris c. crucifer</i>	2	2.74
Ranidae	bullfrog	<i>Rana catesbeiana</i>	14	19.18
	green frog	<i>Rana clamitans melanota</i>	13	17.81
	pickerel frog	<i>Rana palustris</i>	5	6.85
Chelydridae	common snapping turtle	<i>Chelydra serpentina</i>	9	12.33
Emydidae	eastern box turtle	<i>Terrapene c. carolina</i>	7	9.59
	eastern painted turtle	<i>Chrysemys p. picta</i>	1	1.37
	spotted turtle	<i>Clemmys guttata</i>	1	1.37
Colubridae	eastern garter snake	<i>Thamnophis s. sirtalis</i>	1	1.37
	northern water snake	<i>Nerodia s. sipedon</i>	4	5.48
None			17	23.29

Table 5. Physical habitat data for Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

	Instream Habitat ¹	Velocity/Depth Diversity ¹			Riffle Quality ¹	Percent Shading ¹		Number of Woody Debris		Percent Channel Flow ¹		Bank Stability ¹		Aesthetic Rating ¹	
Site		Epifaunal Substrate ¹		Pool Quality ¹		Percent Embeddedness ¹		Maximum Depth (cm) ¹		Number of Rootwads		Channel Alteration ¹		Riparian Width (m) ¹	
CR-P-021-329-96	14	17	10	18	7	17	90	66	2	3	100	5	15	0	19
FR-B-032-206-96	16	5	10	15	16	40	75	27	2	0	97	15	19	10	2
FR-B-046-127-96	6	13	7	6	10	60	98	16	4	0	95	15	17	0	15
FR-B-065-111-96	6	5	8	7	13	45	70	26	2	0	96	7	15	0	6
FR-B-076-118-96	11	4	9	9	8	35	92	24	2	0	94	7	18	5	15
FR-B-081-229-96	18	18	10	15	17	35	85	42	3	2	95	17	16	19	16
FR-B-133-222-96	16	17	10	13	15	25	90	46	2	4	94	15	19	50	18
FR-B-164-137-96	11	4	9	10	10	40	95	30	1	0	70	15	17	50	20
FR-P-005-141-96	17	7	13	15	5	100	92	81	0	0	95	11	19	14	5
FR-P-009-341-96	15	13	13	15	14	20	45	73	2	0	85	12	16	3	6
FR-P-009-347-96	13	10	13	15	13	20	55	74	0	1	80	11	8	0	6
FR-P-015-304-96	15	15	18	19	14	20	45	135	5	4	96	10	5	0	12
FR-P-034-228-96	16	16	12	12	16	73	85	52	0	1	98	8	17	10	11
FR-P-038-139-96	16	16	10	10	17	35	75	37	0	0	98	19	17	6	15
FR-P-046-227-96	12	12	13	15	13	40	20	50	0	0	80	12	6	0	16
FR-P-050-354-96	13	12	17	17	17	20	15	102	1	1	98	6	6	0	12
FR-P-093-237-96	14	5	9	11	11	60	35	36	2	0	100	19	18	0	10
FR-P-093-238-96	13	4	8	17	5	60	20	42	0	0	98	5	18	0	10
FR-P-100-117-96	9	5	9	10	11	60	95	38	3	1	96	8	16	0	10
FR-P-101-233-96	13	15	14	11	18	30	35	58	0	0	97	10	3	0	15
FR-P-103-230-96	6	6	7	8	11	50	93	38	4	1	85	7	8	0	7
FR-P-111-134-96	10	6	6	16	8	90	10	41	0	0	90	5	4	0	11
FR-P-116-221-96	8	12	9	14	11	40	10	48	1	1	95	10	5	0	3
FR-P-132-320-96	11	11	12	16	16	50	82	58	2	0	96	15	10	0	13
FR-P-156-217-96	10	14	11	11	16	25	50	58	5	2	95	10	12	50	16
FR-P-156-231-96	16	16	16	14	15	25	80	82	0	2	95	14	10	0	16
FR-P-156-234-96	18	16	16	18	15	15	70	138	15	4	95	9	13	0	15
FR-P-156-252-96	16	16	16	16	15	25	85	96	6	1	95	10	10	50	16
FR-P-168-218-96	11	7	9	16	13	100	92	38	1	2	97	6	10	3	9
FR-P-214-303-96	19	15	14	19	16	55	70	106	2	2	95	12	9	50	18
FR-P-214-342-96	15	4	16	15	12	60	65	87	1	2	99	6	8	50	16
FR-P-223-225-96	14	16	18	18	16	12	70	108	1	0	95	17	7	0	18
FR-P-223-240-96	9	8	12	13	11	40	45	56	0	0	95	10	8	0	15
FR-P-258-202-96	8	3	17	17	12	80	95	106	3	5	100	19	4	12	14
FR-P-258-215-96	15	13	16	16	16	100	90	59	0	2	100	18	15	7	16
FR-P-258-243-96	14	3	17	18	15	75	75	109	3	1	97	6	6	0	15

Table 5 (cont.). Physical habitat data for Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

	Instream Habitat ¹	Velocity/Depth Diversity ¹		Riffle Quality ¹	Percent Shading ¹		Number of Woody Debris		Percent Channel Flow ¹		Bank Stability ¹		Aesthetic Rating ¹		
Site	Epifaunal Substrate ¹		Pool Quality ¹		Percent Embeddedness ¹		Maximum Depth (cm) ¹		Number of Rootwads		Channel Alteration ¹		Riparian Width (m) ¹		
FR-P-261-122-96	3	2	6	16	11	65	50	29	0	0	95	18	11	0	5
FR-P-263-311-96	12	4	12	16	16	50	40	90	3	1	95	11	12	5	11
FR-P-265-335-96	18	18	16	18	17	30	80	89	7	16	98	16	16	50	12
FR-P-265-351-96	19	18	18	18	19	25	80	72	1	4	98	15	16	50	12
FR-P-275-239-96	16	18	16	12	17	20	70	89	2	3	98	13	10	0	15
FR-P-277-115-96	9	3	7	11	8	65	90	36	3	2	96	5	7	0	16
FR-P-288-133-96	12	4	14	16	14	55	97	54	4	0	97	5	16	50	5
FR-P-290-121-96	14	1	7	5	2	100	95	49	5	1	95	3	17	0	15
FR-P-294-313-96	17	1	7	17	11	50	65	74	1	2	100	20	11	2	13
FR-P-294-357-96	17	9	15	16	9	68	78	64	2	4	99	7	9	33	12
FR-P-298-308-96	17	17	16	15	17	40	75	76	0	0	100	18	13	3	9
FR-P-300-130-96	7	6	9	6	5	100	95	41	0	1	60	6	9	2	12
FR-P-302-334-96	14	16	16	16	14	60	82	79	0	2	90	10	17	0	14
FR-P-319-352-96	16	11	16	17	17	35	30	100	0	1	100	20	10	0	14
FR-P-321-214-96	17	17	17	17	18	19	96	90	1	4	99	16	5	45	13
FR-P-335-110-96	10	1	7	16	4	80	40	56	0	0	100	20	18	8	3
FR-P-349-204-96	12	11	9	11	12	55	64	42	1	3	100	16	5	0	12
FR-P-351-112-96	15	13	11	17	6	80	75	85	0	3	95	10	13	0	16
FR-P-354-321-96	10	2	16	18	12	80	96	102	7	3	100	20	4	11	12
FR-P-360-220-96	13	5	12	15	12	45	85	79	0	0	90	7	6	50	11
FR-P-371-132-96	8	8	8	13	8	70	20	33	1	0	95	10	5	0	15
FR-P-377-242-96	14	12	16	15	13	31	60	95	6	4	75	6	6	50	10
FR-P-388-208-96	15	16	15	14	13	50	65	85	1	0	80	11	6	0	16
FR-P-388-246-96	11	17	9	11	14	15	80	39	1	0	95	11	8	24	15
FR-P-394-317-96	16	14	17	17	16	25	10	122	1	2	100	8	6	0	14
FR-P-399-126-96	8	11	8	11	13	50	68	39	1	0	95	8	5	10	13
FR-P-409-210-96	7	3	11	7	7	55	10	72	0	1	80	9	8	0	16
FR-P-411-305-96	16	16	16	15	17	15	75	88	0	0	95	16	15	0	11
FR-P-421-306-96	19	13	18	19	17	55	90	120	4	2	95	6	13	0	6
FR-P-429-307-96	18	8	13	18	9	60	55	88	1	4	100	12	15	0	6
FR-P-461-251-96	9	10	14	17	11	35	20	80	0	0	98	15	6	0	15
FR-P-462-346-96	9	8	11	13	12	35	30	51	0	0	98	16	11	50	16
FR-P-474-302-96	11	11	18	17	18	65	35	85	3	1	99	16	7	0	16
FR-P-479-348-96	16	19	16	14	18	30	45	70	0	1	96	8	18	8	6
FR-P-516-235-96	15	1	12	15	12	65	25	50	0	2	100	17	5	0	17
FR-P-545-325-96	18	18	18	18	17	15	80	105	10	7	72	8	10	5	16

Table 5 (cont.). Physical habitat data for Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Site	Instream Habitat ¹	Velocity/Depth Diversity ¹		Riffle Quality ¹	Percent Shading ¹		Number of Woody Debris		Percent Channel Flow ¹		Bank Stability ¹	Aesthetic Rating ¹			
	Epifaunal Substrate ¹		Pool Quality ¹		Percent Embeddedness ¹	Maximum Depth (cm) ¹		Number of Rootwads		Channel Alteration ¹		Riparian Width (m) ¹			
FR-P-545-345-96	16	16	16	14	18	15	86	83	5	1	95	16	10	10	18

¹ MBSS Qualitative Habitat Metric - See Appendix B for Guidance




Figure 5. Stream ecological conditions based on the Physical Habitat Index (PHI) at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Table 6. Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Site	Stream Name	F-IBI	B-IBI	PHI
CR-P-021-329-96	Sam's Cr	2.71	1.0	61.31
FR-B-032-206-96	Friends Cr	3.00	1.9	43.23
FR-B-046-127-96	Un Trib To Little Catoctin Cr	1.00	3.2	8.95
FR-B-065-111-96	Un Trib To Catoctin Cr		3.4	11.78
FR-B-076-118-96	Un Trib To Little Catoctin Cr		1.7	19.80
FR-B-081-229-96	Hunting Cr	2.71	2.1	81.28
FR-B-085-212-96	Catoctin Cr		1.4	
FR-B-133-222-96	Owens Cr	4.14	2.1	87.20
FR-B-164-137-96	Middle Cr	2.71	2.1	26.70
FR-P-005-141-96	Un Trib To Carroll Cr	1.00	1.4	14.84
FR-P-009-341-96	Flat Run	3.86	1.4	53.42
FR-P-009-347-96	Flat Run	3.57	1.0	53.42
FR-P-015-304-96	Bennet Cr	4.71	2.3	92.21
FR-P-034-228-96	Little Hunting Cr	3.29	3.0	57.57
FR-P-038-139-96	Turkey Cr	1.00	2.6	60.93
FR-P-046-227-96	Haines Br	2.14	1.7	45.25
FR-P-050-354-96	Cabbage Run	2.71	2.1	79.67
FR-P-093-237-96	Tuscarora Cr	1.57	2.6	23.62
FR-P-093-238-96	Tuscarora Cr	1.57	2.6	10.37
FR-P-100-117-96	Un Trib To Ballenger Cr	2.43	3.9	19.48
FR-P-101-233-96	Fahrney Br	3.86	3.0	67.95
FR-P-103-230-96	Ballenger Cr	3.00	2.8	11.57
FR-P-111-134-96	Un Trib To Piney Cr	2.43	1.0	6.12
FR-P-116-221-96	Un Trib To Israel Cr	3.29	2.3	17.04
FR-P-132-320-96	Little Hunting Cr	4.71	3.7	42.23
FR-P-156-217-96	South Fork Linganore Cr	3.86	2.6	64.75
FR-P-156-231-96	South Fork Linganore Cr	4.43	3.7	86.26
FR-P-156-234-96	South Fork Linganore Cr	4.14	2.8	94.79
FR-P-156-252-96	South Fork Linganore	4.14	3.2	81.28
FR-P-168-218-96	Tuscarora Cr	2.14	1.2	25.13
FR-P-214-303-96	Owens Cr	4.43	2.1	85.76
FR-P-214-342-96	Owens Cr	4.71	1.7	71.41
FR-P-223-225-96	Beaver Dam Cr	2.43	2.6	81.21
FR-P-223-240-96	Beaver Dam Br	1.57	1.4	27.51
FR-P-258-202-96	Fishing Cr	3.57	2.6	72.23
FR-P-258-215-96	Fishing Cr	4.14	3.0	71.41
FR-P-258-243-96	Fishing Cr	3.57	1.9	66.59
FR-P-261-122-96	Un Trib To Hunting Cr	3.29	1.2	4.07
FR-P-263-311-96	Tuscarora Cr	2.71	1.9	52.40
FR-P-265-335-96	Middle Cr	4.14	1.4	99.93
FR-P-265-351-96	Middle Cr	4.43	2.1	97.11
FR-P-275-239-96	Church Br Of Bush Cr	4.71	2.6	92.21
FR-P-277-115-96	Un Trib To Ballenger Cr	2.14	2.6	18.53
FR-P-288-133-96	Trib To Hunting Cr	4.14	3.2	35.90
FR-P-290-121-96	Un Trib To Tuscarora Cr	1.57	1.7	7.99
FR-P-294-313-96	Hunting Cr	3.57	1.9	48.31
FR-P-294-357-96	Hunting Cr	3.86	2.6	75.85
FR-P-298-308-96	Owens Cr	4.14	2.8	73.05
FR-P-300-130-96	Un Trib To Monocacy R	1.29	1.0	5.67

Table 6 (cont.). Fish Index of Biotic Integrity (F-IBI), Benthic Macroinvertebrate Index of Biotic Integrity (B-IBI), and Physical Habitat Index (PHI) scores at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Site	Stream Name	F-IBI	B-IBI	PHI
FR-P-302-334-96	Owens Cr	4.43	1.7	72.23
FR-P-319-352-96	Owens Cr	5.00	2.6	81.89
FR-P-321-214-96	Woodville Br	4.43	2.3	95.85
FR-P-335-110-96	Un Trib To Monocacy R	2.43	1.4	3.55
FR-P-349-204-96	Ballenger Cr	4.43	2.3	48.82
FR-P-351-112-96	Un Trib To Fahrney Br		3.2	42.23
FR-P-354-321-96	Tuscarora Cr	3.00	2.8	56.46
FR-P-360-220-96	Un Trib To Bush Cr	3.29	3.2	35.90
FR-P-371-132-96	Motter's Run	1.57	1.2	8.95
FR-P-377-242-96	Bennett Cr	3.86	2.3	85.36
FR-P-388-208-96	Talbot Br	3.57	1.7	57.46
FR-P-388-246-96	Talbot Br	4.14	2.3	39.75
FR-P-394-317-96	Cabbage Run	3.00	1.9	88.08
FR-P-399-126-96	Un Trib To Weldon Cr		1.7	17.62
FR-P-409-210-96	Addison Run	1.57	2.6	16.75
FR-P-411-305-96	Bens Br	4.14	3.7	77.96
FR-P-421-306-96	Bush Cr	4.14	3.0	87.20
FR-P-429-307-96	Ballenger Cr	3.86	3.7	70.56
FR-P-461-251-96	Ben's Br	3.86	3.2	34.04
FR-P-462-346-96	Clemson Br	1.29	2.3	29.60
FR-P-474-302-96	Sam's Cr	1.57	1.7	72.64
FR-P-479-348-96	Owens Cr	3.86	2.1	79.67
FR-P-516-235-96	Sandy Run	3.29	2.8	59.95
FR-P-545-325-96	Bush Cr	4.43	2.8	98.94
FR-P-545-345-96	Bush Cr	4.43	1.7	88.51

Table 7. Water chemistry data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Site	pH	Conductivity ($\mu\text{S}/\text{cm}$)	Acid Neutralizing Capacity ($\mu\text{eq}/\text{L}$)	Nitrate (mg/L)	Sulfate (mg/L)	Dissolved Oxygen (mg/L)	Dissolved Organic Carbon (mg/L)
FR-B-032-206-96	7.48	0.248	675.90	1.814	17.676	8.50	1.20
FR-B-046-127-96	6.98	0.097	360.20	0.403	14.932	8.20	0.70
FR-B-065-111-96	7.21	0.360	831.30	10.304	37.060	7.80	1.20
FR-B-076-118-96	7.39	0.343	624.00	3.319	23.082	5.50	1.90
FR-B-081-229-96	7.22	0.133	284.00	1.113	6.720	9.50	1.30
FR-B-085-212-96	7.20	0.156	427.30	1.791	15.429		1.70
FR-B-133-222-96	7.33	0.113	324.90	1.347	6.645	9.80	1.30
FR-B-164-137-96	7.07	0.128	274.00	0.734	9.542	8.10	2.00
FR-P-005-141-96	7.28	0.359	790.30	1.129	21.498	11.50	1.60
FR-P-009-341-96	7.49	0.202	1080.00	0.430	27.174	8.60	4.20
FR-P-009-347-96	7.63	0.198	1051.90	0.397	26.258	9.60	3.80
FR-P-015-304-96	7.13	0.153	445.80	3.713	7.492	9.30	1.40
FR-P-034-228-96	6.86	0.057	131.90	0.428	6.292	8.70	1.20
FR-P-038-139-96	7.22	0.089	312.60	0.572	7.436	9.30	1.30
FR-P-046-227-96	7.77	0.439	2667.00	7.706	26.444	8.60	2.00
FR-P-050-354-96	7.61	0.417	1070.30	2.280	114.454	8.60	2.10
FR-P-093-237-96	8.05	0.217	1109.10	0.960	13.017	9.50	5.80
FR-P-093-238-96	7.88	0.216	1104.20	0.954	13.056	9.50	5.90
FR-P-100-117-96	7.23	0.190	440.10	2.389	15.122	8.80	1.20
FR-P-101-233-96	7.37	0.168	516.40	4.275	10.723	9.90	2.80
FR-P-103-230-96	8.55	0.268	1051.20	4.543	17.807	10.10	1.80
FR-P-111-134-96	7.08	0.255	526.60	7.959	28.006	8.60	3.60
FR-P-116-221-96	7.24	0.145	345.60	2.708	9.209	9.90	1.80
FR-P-132-320-96	7.25	0.084	311.20	0.514	7.406	9.50	1.90
FR-P-156-217-96	7.34	0.115	271.10	3.102	4.326	9.90	1.10
FR-P-156-231-96	7.22	0.111	265.60	3.042	4.339	10.10	1.30
FR-P-156-234-96	7.29	0.112	274.30	3.014	4.510	9.20	0.90
FR-P-156-252-96	7.07	0.113	271.10	2.616	5.451	10.00	2.00
FR-P-168-218-96	7.72	0.401	1662.70	5.507	25.856	9.00	3.20
FR-P-214-303-96	7.81	0.146	554.70	1.377	12.511	7.60	1.70
FR-P-214-342-96	7.62	0.148	556.70	1.378	11.249	8.70	1.70
FR-P-223-225-96	7.62	0.305	1831.30	4.708	16.240	9.70	2.90
FR-P-223-240-96	7.75	0.296	1440.20	5.442	16.528	9.20	1.60
FR-P-258-202-96	7.57	0.153	514.80	0.937	10.008	8.10	5.00
FR-P-258-215-96	7.44	0.125	436.50	0.808	9.209	8.90	4.00
FR-P-258-243-96	7.28	0.073	237.80	0.725	7.101	7.60	1.90
FR-P-261-122-96	7.28	0.129	353.10	3.589	8.688	8.10	2.20
FR-P-263-311-96	7.75	0.357	1892.60	5.454	24.755	9.30	2.00
FR-P-265-335-96	8.75	0.168	1034.30	0.944	14.815	9.00	2.70
FR-P-265-351-96	8.65	0.164	1041.80	0.989	14.904	9.30	2.50
FR-P-275-239-96	7.25	0.139	315.30	3.377	5.534	9.60	1.10
FR-P-277-115-96	7.56	0.282	743.30	4.087	19.087	7.50	2.00
FR-P-288-133-96	7.33	0.120	292.80	0.559	6.487	8.70	1.70
FR-P-290-121-96	7.18	0.278	1899.90	4.573	17.645	7.60	1.50
FR-P-294-313-96	7.61	0.175	461.10	0.946	8.605	7.80	1.90
FR-P-294-357-96	7.49	0.197	583.10	0.803	8.603	7.60	1.10
FR-P-298-308-96	7.50	0.141	517.30	1.453	10.431	7.60	1.60
FR-P-300-130-96	7.55	0.282	770.40	5.981	33.303	5.20	5.20

Table 7 (cont.). Water chemistry data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

Site	pH	Conductivity (μ S/cm)	Acid Neutralizing Capacity (μ eq/L)	Nitrate (mg/L)	Sulfate (mg/L)	Dissolved Oxygen (mg/L)	Dissolved Organic Carbon (mg/L)
FR-P-302-334-96	7.49	0.124	370.20	1.016	9.065	8.60	1.50
FR-P-319-352-96	7.63	0.147	551.20	1.269	9.598	9.30	1.40
FR-P-321-214-96	7.19	0.138	438.10	4.075	6.645	8.60	1.00
FR-P-335-110-96	7.70	0.707	4148.30	2.789	30.740	5.80	1.90
FR-P-349-204-96	7.88	0.333	1592.00	4.158	17.358	9.20	1.40
FR-P-351-112-96	7.05	0.145	467.50	3.540	8.615	7.50	1.90
FR-P-354-321-96	7.73	0.385	2151.40	6.134	25.168	7.80	2.20
FR-P-360-220-96	8.05	0.237	880.50	5.542	8.056	7.60	2.30
FR-P-371-132-96	7.57	0.286	1087.10	2.704	24.896	6.30	3.90
FR-P-377-242-96	7.17	0.125	264.30	3.026	7.221	9.20	1.70
FR-P-388-208-96	7.24	0.153	301.60	2.829	5.411	10.20	0.80
FR-P-388-246-96	7.43	0.179	303.30	3.117	5.497	8.50	0.90
FR-P-394-317-96	7.73	0.433	976.70	2.270	125.827	8.50	1.50
FR-P-399-126-96	6.77	0.185	511.60	7.096	7.279	9.90	1.70
FR-P-409-210-96	7.92	0.292	1519.40	3.849	18.209	11.60	2.90
FR-P-411-305-96	7.64	0.197	1141.00	3.681	8.333	8.90	1.20
FR-P-421-306-96	8.74	0.244	739.00	2.932	11.311	8.30	1.90
FR-P-429-307-96	8.30	0.369	1825.90	4.565	22.088	9.60	1.80
FR-P-461-251-96	7.38	0.139	631.00	2.492	8.364	9.40	2.60
FR-P-462-346-96	7.88	0.324	1902.80	6.157	14.463	11.50	1.40
FR-P-474-302-96	8.32	0.345	1948.70	5.432	24.018	9.00	2.10
FR-P-479-348-96	7.52	0.132	388.90	1.238	10.089	8.40	1.60
FR-P-516-235-96	7.81	0.243	1561.20	1.850	11.395	7.50	1.70
FR-P-545-325-96	8.77	0.251	690.10	2.891	10.171	9.60	1.60
FR-P-545-345-96	8.61	0.249	688.50	2.917	10.135	9.30	1.40




Figure 6. Nitrate-nitrogen concentrations (mg/L) at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Literature Cited

- Barbour, M.T., and J.B. Stribling. 1991. Use of habitat assessment in evaluating the biological integrity of stream communities. In: *Biological Criteria: Research and Regulation*. U.S. Environmental Protection Agency, Washington, D.C. EPA-440/5-91-005. pp. 25-38.
- Cairns, J. and J.R. Pratt. 1993. A history of biological monitoring using benthic macroinvertebrates. In Rosenberg, D.M. and V.H. Resh, eds. *Freshwater Monitoring and Benthic Macroinvertebrates*. Chapman and Hall, New York
- CBRM (Chesapeake Bay Research and Monitoring Division). 1989. Acid deposition in Maryland. Summary of results through 1988. Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, Annapolis, MD. AD-89-1. NTIS No. PB89-182729/AS.
- Chaillou, J.C. 1995. Technical Memorandum, The Maryland Biological Stream Survey: Summary of landowner contact procedures. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division, Annapolis, MD. CBWP-MANTA-EA-95-1.
- Cochran, W.G. 1977. *Sampling Techniques*. 3rd ed. New York: John Wiley and Sons.
- EPA (U.S. Environmental Protection Agency). 1987. *Handbook of methods for acid deposition studies: Laboratory analysis for surface water chemistry*. EPA-600/4-87/026.
- Hall, L.W., Jr., R.P. Morgan II, E.S. Perry, A. Waltz. 1999. Development of a provisional physical habitat index for Maryland freshwater streams. Prepared by Wye Research and Education Center, Queenstown, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment 455.
- Heimbuch, D.G., J.C. Seibel, H.T. Wilson, P.F. Kazyak. 1999. A multiyear lattice sampling design for Maryland-wide fish abundance estimation. *Journal of Agricultural, Biological, and Environmental Statistics*. 4: 443-455.
- Hilsenhoff, W.L. 1987. An improved biotic index of organic stream pollution. *Great Lakes Entomologist* 20: 31-39.
- Janicki, A., D. Wade, D. Heimbuch, H. Wilson, P. Jacobson, P. Kazyak. 1993. Maryland Biological Stream Survey design report. Prepared for Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, by Coastal Environmental Services, Linthicum, MD. CBRM-AD-93-1.
- Jessen, R.J. 1978. *Statistical Survey Techniques*. John Wiley, New York.
- Karr, J.R. 1991. Biological integrity: A long-neglected aspect of water resource management. *Ecological Applications* 1: 66-84.
- Karr, J.R., K.D. Fausch, P.L. Angermeier, P.R. Yant, and I.J. Schlosser. 1986. Assessing biological integrity in running waters: a method and its rationale. *Illinois Natural History Survey Special Publication* 5. 28 pp.
- Kazyak, P.F. 1996. Maryland Biological Stream Survey Sampling Manual. Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division.
- Kazyak, P.F. 1994. Maryland Biological Stream Survey Sampling Manual. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division.
- Kerans, B.L. and J.R. Karr. 1994. A benthic index of biotic integrity (B-IBI) for rivers of the Tennessee valley. *Ecological Applications* 4: 768-785.
- Knapp, C.M., W.P. Saunders, D.G. Heimbuch, H.S. Greening, and G.J. Filbin. 1988. Maryland

- Synoptic Stream Chemistry Survey: Estimating the number and distribution of streams affected by or at risk from acidification. Prepared by International Science and Technology, Inc., Reston, VA, for the Maryland Department of Natural Resources, Power Plant Research Program, Annapolis, MD. AD-88-2. NTIS No. PB88-213996/AS.
- Knapp, C.M. and W.P. Saunders. 1987. Maryland Synoptic Stream Chemistry Survey design report. Prepared by International Science and Technology, Inc. for Maryland Department of Natural Resources, Power Plant Research Program. AD-87-5.
- Lenat, D.R. 1988. Water quality assessment of streams using a qualitative collection method for benthic macroinvertebrates. *Journal of North American Benthological Society* 7: 222-223.
- Miller, D.L., P.M. Leonard, R.M. Hughes, J.R. Karr, P.B. Moyle, L.H. Schrader, B.A. Thompson, R.A. Daniels, K.S. Fausch, G.A. Fitzhugh, J.R. Gammon, D.B. Halliwell, P.L. Angermeier, and D.J. Orth. 1988. Regional applications of an index of biotic integrity for use in water resource management. *Fisheries* 13(5): 12-20.
- Ohio EPA (Ohio Environmental Protection Agency). 1987. Biological criteria for the protection of aquatic life. Volumes I-III. Ohio EPA, Division of Water Quality Monitoring and Assessment, Surface Water Section, Columbus, Ohio.
- Plafkin, J.L., M.T. Barbour, K.D. Porter, S.K. Gross and R.M. Hughes. 1989. Rapid bioassessment protocols for use in streams and rivers: Benthic macroinvertebrates and fish. U.S. Environmental Protection Agency, Office of Water, Washington, D.C. EPA 440-4-89-001.
- Ranasinghe, J.A., S.B. Weisberg, D.M. Dauer, L.C. Schaffner, R.J. Diaz, and J.B. Frithsen. 1994. Chesapeake Bay Benthic Community Restoration Goals. Prepared by Versar, Inc. for U.S. EPA and the Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division.
- Rankin, E.T. 1989. The Qualitative Habitat Evaluation Index (QHEI): Rationale, methods, and application. Ohio EPA, Division of Water Quality Planning and Assessment, Ecological Analysis Section, Columbus, OH.
- Resh, V. H., 1995. Freshwater benthic macroinvertebrates and rapid assessment procedures for water quality monitoring in the developing and newly industrialized countries. In: Davis, W.S. and T.P. Simon, eds. 1995. Biological assessment and criteria: tools for water resource planning and decision making. Lewis Publishers, Boca Raton, FL.
- Roth, N.E., M.T. Southerland, G. Mercurio, J.C. Chaillou, P.F. Kazyak, S.A. Stranko, A.P. Prochaska, D.G. Heimbuch, and J.C. Seibel. 1999. State of the Streams: 1995-1997 Maryland Biological Stream Survey Results. Prepared by Versar, Inc., Columbia, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division. CBWP-MANTA-EA-99-6.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, G.T. Wilson, D.G. Heimbuch, and J.C. Seibel. 1998a. Maryland Biological Stream Survey: Ecological Status of Non-tidal Streams Sampled in 1996. Prepared by Versar, Inc., Columbia, MD, and Post, Buckley, Schuh, and Jernigan, Inc., Bowie, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division. CBWP-MANTA-EA-97-2.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, R.J. Klauda, P.F. Kazyak, S.A. Stranko, S.B. Weisberg, L.W. Hall, Jr., and R.P. Morgan II. 1998b. Maryland Biological Stream Survey: Development of a Fish Index of Biotic Integrity. *Environmental Management and Assessment* 51: 89-106.
- Roth, N.E., M.T. Southerland, J.C. Chaillou, J.H. Volstad, S.B. Weisberg, H.T. Wilson, D.G. Heimbuch, and J.C. Seibel. 1997. Maryland Biological Stream Survey: Ecological Status of Non-Tidal Streams in Six Basins Sampled in 1995. Prepared by Versar, Inc., Columbia, MD and Coastal Environmental Services, Linthicum, MD for Maryland Department of Natural Resources,

Monitoring and Non-tidal Assessment Division.
CBWP-MANTA-EA-97-2.

Simon, T.P. ed. 1999. Assessing the Sustainability and Biological Integrity of Water Resources Using Fish Communities. CRC Press, Washington DC.

Strahler, A.N. 1957. Quantitative analysis of watershed geomorphology. Transactions of the American Geophysical Union 38(6): 913-920.

Stribling, J.B., B.K. Jessup, J.S. White, D. Boward, M. Hurd. 1998. Development of a benthic index of biotic integrity for Maryland streams. Prepared by Tetra Tech, Inc., Owings Mills, MD, for Maryland Department of Natural Resources, Monitoring and Non-tidal Assessment Division, Annapolis, MD. CBWP-EA-98-3.

Tornatore, T., P. Jacobson, J. Gerritsen and A. Janicki. 1992. Data sets for developing measures of biological integrity in Maryland streams. Prepared by Versar, Inc., Columbia, MD, for the Maryland Department of Natural Resources, Chesapeake Bay Research and Monitoring Division, Annapolis, MD. AD-93-7.

Weisberg, S.B., J.A. Ranasinghe, D.M. Dauer, L.C. Schaffner, R.J. Diaz, J.B. Frithsen. 1997. An estuarine benthic index of biotic integrity (B-IBI) for Chesapeake Bay. Estuaries 20: 149-158.

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Appendix A. Summary of the types of data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997. Abbreviations used are as follows: F-IBI - Fish Index of Biotic Integrity; B-IBI Benthic Macroinvertebrate Index of Biotic Integrity; Fam.IBI - Family-Level Benthic Macroinvertebrate Index of Biotic Integrity; PHI - Physical Habitat Index.

Site	Stream Name	Benthic Macroinvertebrate		Habitat		F-IBI	Fam. IBI	
		Fish	Herpetofauna	Water Chemistry			B-IBI	PHI
CR-P-021-329-96	Sam's Cr	X	X	X	X	X	X	X
FR-B-032-206-96	Friends Cr	X	X	X	X	X	X	X
FR-B-046-127-96	Un Trib To Little Catoctin Cr	X	X	X	X	X	X	X
FR-B-065-111-96	Un Trib To Catoctin Cr	X	X	X	X		X	X
FR-B-076-118-96	Un Trib To Little Catoctin Cr	X	X	X	X		X	X
FR-B-081-229-96	Hunting Cr	X	X	X	X	X	X	X
FR-B-085-212-96	Catoctin Cr		X		X		X	
FR-B-133-222-96	Owens Cr	X	X	X	X	X	X	X
FR-B-164-137-96	Middle Cr	X	X	X	X	X	X	X
FR-P-005-141-96	Un Trib To Carroll Cr	X	X	X	X	X	X	X
FR-P-009-341-96	Flat Run	X	X	X	X	X	X	X
FR-P-009-347-96	Flat Run	X	X	X	X	X	X	X
FR-P-015-304-96	Bennet Cr	X	X	X	X	X	X	X
FR-P-034-228-96	Little Hunting Cr	X	X	X	X	X	X	X
FR-P-038-139-96	Turkey Cr	X	X	X	X	X	X	X
FR-P-046-227-96	Haines Br	X	X	X	X	X	X	X
FR-P-050-354-96	Cabbage Run	X	X	X	X	X	X	X
FR-P-093-237-96	Tuscarora Cr	X	X	X	X	X	X	X
FR-P-093-238-96	Tuscarora Cr	X	X	X	X	X	X	X
FR-P-100-117-96	Un Trib To Ballenger Cr	X	X	X	X	X	X	X
FR-P-101-233-96	Fahrney Br	X	X	X	X	X	X	X
FR-P-103-230-96	Ballenger Cr	X	X	X	X	X	X	X
FR-P-111-134-96	Un Trib To Piney Cr	X	X	X	X	X	X	X
FR-P-116-221-96	Un Trib To Israel Cr	X	X	X	X	X	X	X
FR-P-132-320-96	Little Hunting Cr	X	X	X	X	X	X	X
FR-P-156-217-96	South Fork Linganore Cr	X	X	X	X	X	X	X
FR-P-156-231-96	South Fork Linganore Cr	X	X	X	X	X	X	X
FR-P-156-234-96	South Fork Linganore Cr	X	X	X	X	X	X	X
FR-P-156-252-96	South Fork Linganore	X	X	X	X	X	X	X
FR-P-168-218-96	Tuscarora Cr	X	X	X	X	X	X	X
FR-P-214-303-96	Owens Cr	X	X	X	X	X	X	X
FR-P-214-342-96	Owens Cr	X	X	X	X	X	X	X
FR-P-223-225-96	Beaver Dam Cr	X	X	X	X	X	X	X
FR-P-223-240-96	Beaver Dam Br	X	X	X	X	X	X	X
FR-P-258-202-96	Fishing Cr	X	X	X	X	X	X	X

Appendix A (cont.). Summary of the types of data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997. Abbreviations used are as follows: F-IBI - Fish Index of Biotic Integrity; B-IBI - Benthic Macroinvertebrate Index of Biotic Integrity; Fam. IBI - Family-Level Benthic Macroinvertebrate Index of Biotic Integrity; PHI - Physical Habitat Index.

Site	Stream Name	Benthic Macroinvertebrate		Habitat		F-IBI		Fam. IBI	
		Fish	Herpetofauna	Water Chemistry			B-IBI		PHI
FR-P-258-215-96	Fishing Cr	X	X	X	X	X	X	X	X
FR-P-258-243-96	Fishing Cr	X	X	X	X	X	X	X	X
FR-P-261-122-96	Un Trib To Hunting Cr	X	X	X	X	X	X	X	X
FR-P-263-311-96	Tuscarora Cr	X	X	X	X	X	X	X	X
FR-P-265-335-96	Middle Cr	X	X	X	X	X	X	X	X
FR-P-265-351-96	Middle Cr	X	X	X	X	X	X	X	X
FR-P-275-239-96	Church Br Of Bush Cr	X	X	X	X	X	X	X	X
FR-P-277-115-96	Un Trib To Ballenger Cr	X	X	X	X	X	X	X	X
FR-P-288-133-96	Trib To Hunting Cr	X	X	X	X	X	X	X	X
FR-P-290-121-96	Un Trib To Tuscarora Cr	X	X	X	X	X	X	X	X
FR-P-294-313-96	Hunting Cr	X	X	X	X	X	X	X	X
FR-P-294-357-96	Hunting Cr	X	X	X	X	X	X	X	X
FR-P-298-308-96	Owens Cr	X	X	X	X	X	X	X	X
FR-P-300-130-96	Un Trib To Monocacy R	X	X	X	X	X	X	X	X
FR-P-302-334-96	Owens Cr	X	X	X	X	X	X	X	X
FR-P-319-352-96	Owens Cr	X	X	X	X	X	X	X	X
FR-P-321-214-96	Woodville Br	X	X	X	X	X	X	X	X
FR-P-335-110-96	Un Trib To Monocacy R	X	X	X	X	X	X	X	X
FR-P-349-204-96	Ballenger Cr	X	X	X	X	X	X	X	X
FR-P-351-112-96	Un Trib To Fahrney Br	X	X	X	X		X	X	X
FR-P-354-321-96	Tuscarora Cr	X	X	X	X	X	X	X	X
FR-P-360-220-96	Un Trib To Bush Cr	X	X	X	X	X	X	X	X
FR-P-371-132-96	Motter's Run	X	X	X	X	X	X	X	X
FR-P-377-242-96	Bennett Cr	X	X	X	X	X	X	X	X
FR-P-388-208-96	Talbot Br	X	X	X	X	X	X	X	X
FR-P-388-246-96	Talbot Br	X	X	X	X	X	X	X	X
FR-P-394-317-96	Cabbage Run	X	X	X	X	X	X	X	X
FR-P-399-126-96	Un Trib To Weldon Cr	X	X	X	X		X	X	X
FR-P-409-210-96	Addison Run	X	X	X	X	X	X	X	X
FR-P-411-305-96	Bens Br	X	X	X	X	X	X	X	X
FR-P-421-306-96	Bush Cr	X	X	X	X	X	X	X	X
FR-P-429-307-96	Ballenger Cr	X	X	X	X	X	X	X	X
FR-P-461-251-96	Ben's Br	X	X	X	X	X	X	X	X
FR-P-462-346-96	Clemson Br	X	X	X	X	X	X	X	X
FR-P-474-302-96	Sam's Cr	X	X	X	X	X	X	X	X

Appendix A (cont.) Summary of the types of data collected at Maryland Biological Stream Survey sites in Frederick County, 1994-1997.
 Abbreviations used are as follows: F-IBI - Fish Index of Biotic Integrity; B-IBI - Benthic Macroinvertebrate Index of Biotic Integrity; Fam. IBI - Family-Level Benthic Macroinvertebrate Index of Biotic Integrity; PHI - Physical Habitat Index.

Site	Stream Name	Benthic Macroinvertebrate		Habitat		F-IBI		Fam. IBI	PHI
		Fish	Herpetofauna	Water Chemistry			B-IBI		
FR-P-479-348-96	Owens Cr	X	X	X	X	X	X	X	X
FR-P-516-235-96	Sandy Run	X	X	X	X	X	X	X	X
FR-P-545-325-96	Bush Cr	X	X	X	X	X	X	X	X
FR-P-545-345-96	Bush Cr	X	X	X	X	X	X	X	X

**THIS PAGE INTENTIONALLY
LEFT BLANK**

Appendix B. Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997. All variables rated on a scale of 0 (poor) to 20 (optimal) unless otherwise noted.

SUBSTRATE AND INSTREAM COVER

Instream Habitat is rated according to the perceived value of habitat to the fish community. Higher scores are assigned to sites with a variety of habitat types and particle sizes. In addition, higher scores are assigned to sites with a high degree of uneven substrate, including logs and rootwads. In streams where substrate types are favorable but flows are so low that fish are essentially precluded from using the habitat, low scores are assigned. If none of the habitat within a segment is useable by fish, a score of zero is assigned.

Epifaunal Substrate is rated based on the amount and variety of hard, stable substrates usable by benthic macroinvertebrates. Because they inhibit colonization, flocculent materials or fine sediments surrounding otherwise good substrates are assigned low scores. Scores are also reduced when substrates are less stable.

Velocity/Depth Diversity is rated based on the variety of velocity/depth regimes present at a site (slow-shallow, slow-deep, fast-shallow, and fast-deep). As with embeddedness, this metric varies by stream gradient.

Pool/Glide/Eddy Quality is rated based on the variety and spatial complexity of slow or still water habitat within the sample segment. In high-gradient streams, functionally important slow water habitat may exist in the form of larger eddies. Within a category, higher scores are assigned to segments which have undercut banks, woody debris or other types of cover for fish.

Riffle/Run Quality is based on the depth, complexity, and functional importance of riffle/run habitat in the segment, with highest scores assigned to segments dominated by deeper riffle/run areas, stable substrates, and a variety of current velocities.

Embeddedness is a percentage of surface area of larger particles that is surrounded by fine sediments on the stream bottom. In low gradient streams, embeddedness may be high even in relatively unimpaired watersheds.

CHANNEL CHARACTER

Channel Alteration is a measure of large-scale changes in the shape of the stream channel. Channel alteration includes: concrete channels, artificial embankments, obvious straightening of the natural channel, rip-rap, or other structures, as well as recent bar development. Ratings for this metric are based on the presence of artificial structures as well as the existence, extent, and coarseness of point bars, side bars, and mid-channel bars which indicate the degree of flow fluctuations and substrate stability. Evidence of channelization may sometimes be seen in the form of berms that parallel the stream channel.

Bank Stability is rated based on the presence/absence of riparian vegetation and other stabilizing bank materials such as boulders and rootwads, and frequency/size of erosional areas. Sites with steep slopes are not penalized if banks are composed solely of stable materials.

Channel Flow Status is the percentage of the stream channel that has water, with subtractions made for exposed substrates and dewatered areas.

RIPARIAN CORRIDOR

Shading is rated based on estimates of the degree and duration of shading at a site during summer, including any effects of shading caused by land forms.

Appendix B (cont.).Physical habitat condition measured by the Maryland Biological Stream Survey, 1994-1997.
All variables rated on a scale of 0 (poor) to 20 (optimal) unless otherwise noted.

Riparian Buffer is rated according to the size and type of the vegetated riparian buffer zone at the site. Cultivated fields for agriculture that have bare soil to any extent are not considered as riparian buffers. At sites where the buffer width is variable, or direct delivery of storm runoff or sediment to the stream is evident or highly likely, the narrowest representative buffer width in the segment (e.g., 0 if parking lot runoff enters directly to the stream) is measured and recorded even though some of the stream segment may have a well developed riparian buffer.

AESTHETICS/REMOTENESS

Aesthetics are rated according to the visual appeal of the site and presence/absence of human refuse, with highest scores assigned to stream segments with no human refuse and visually outstanding character.

Remoteness is rated based on the absence of detectable human activity and difficulty in accessing the segment.